

Multiscale ecology of agroecosystems is an emerging research field that can provide a stronger theoretical background for the integrated pest management

Petrovskii, Sergei; Petrovskaya, Natalia; Bearup, Daniel

DOI:

[10.1016/j.plrev.2014.07.001](https://doi.org/10.1016/j.plrev.2014.07.001)

License:

Other (please specify with Rights Statement)

Document Version

Peer reviewed version

Citation for published version (Harvard):

Petrovskii, S, Petrovskaya, N & Bearup, D 2014, 'Multiscale ecology of agroecosystems is an emerging research field that can provide a stronger theoretical background for the integrated pest management', *Physics of Life Reviews*, vol. 11, no. 3, pp. 536-539. <https://doi.org/10.1016/j.plrev.2014.07.001>

[Link to publication on Research at Birmingham portal](#)

Publisher Rights Statement:

NOTICE: this is the author's version of a work that was accepted for publication in *Physics of Life Reviews*. Changes resulting from the publishing process, such as peer review, editing, corrections, structural formatting, and other quality control mechanisms may not be reflected in this document. Changes may have been made to this work since it was submitted for publication. A definitive version was subsequently published in *Physics of Life Reviews*, Volume 11, Issue 3, September 2014, Pages 536–539 DOI: 10.1016/j.plrev.2014.07.001
Checked for repository 28/10/2014

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Accepted Manuscript


Multiscale ecology of agroecosystems is an emerging research field that can provide a stronger theoretical background for the integrated pest management

Sergei Petrovskii, Natalia Petrovskaya, Daniel Bearup

PII: S1571-0645(14)00114-6
DOI: [10.1016/j.plrev.2014.07.001](http://dx.doi.org/10.1016/j.plrev.2014.07.001)
Reference: PLREV 521

To appear in: *Physics of Life Reviews*

Received date: 30 June 2014
Accepted date: 1 July 2014



Please cite this article in press as: Petrovskii S, et al. Multiscale ecology of agroecosystems is an emerging research field that can provide a stronger theoretical background for the integrated pest management. *Phys Life Rev* (2014), <http://dx.doi.org/10.1016/j.plrev.2014.07.001>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Reply to comment

Multiscale ecology of agroecosystems is an emerging research field that can provide a stronger theoretical background for the integrated pest management

Reply to comments on “Multiscale approach to pest insect monitoring: Random walks, pattern formation, synchronization, and networks”

By Sergei Petrovskii¹, Natalia Petrovskaya², and Daniel Bearup³

¹ Department of Mathematics, University of Leicester, Leicester, LE1 7RH, U.K.

² School of Mathematics, University of Birmingham, Birmingham, B15 2TT, U.K.

³ ICBM, University of Oldenburg, 26111 Oldenburg, Germany

Keywords: Insect pest monitoring; Baited trap; Pitfall trap; Taxis; Random walk; Lévy flight

We would like to thank all commentators for their insightful and thought-provoking commentaries [1-4] that also helped to further broaden the scope of our review [5] as well as to extend the list of references. We very much appreciate the positive comments on the relevance, timeliness and comprehensiveness of our work.

Meanwhile, we want to emphasize that multiscale pest monitoring is an emerging research theme rather than a well-established one. Indeed, although the existence and importance of multiple scales has long been known in spatial ecology [6,7], their quantitative investigation has not started in full until relatively recently, e.g. see [8,9] and references therein. For this reason, in our review [5], several relevant aspects might have been either just touched briefly or not mentioned at all. In particular, Rebecca Tyson [1] rightfully brought attention to the importance of transient environmental factors, e.g. temperature, that may control the build-up of pest populations. The effects of temperature on the rates of insect development through their life-stages and on insect population growth are well-known [10,11] and have been utilized in some empirical approaches used in Integrated Pest Management (IPM), e.g. in what is known as degree-day models [12,13]. Models of this type, however, are essentially based on average insect population numbers and hence leave largely unaddressed the natural spatial and temporal variability of the insect population between subsequent years and between spatial locations (e.g. agricultural fields) with similar properties. Meanwhile, such variability is typical both on the field and landscape scales, and can often be as large as two orders of magnitude, e.g. see Figs.23 and 32 in [5] and Fig.2 in [14]. Multiscale pest monitoring can provide a stronger theoretical background for IPM and to transform empirical models to theory-based models, and, at the same time, take a full account of population variability. The currently available theoretical approaches to the analysis, interpretation and modelling of trap counts (in particular those reviewed in [5]) are relatively straightforward to extend to the fully realistic case of a transient environment, provided sufficient information is available about insect response to all relevant factors. The latter may require additional field studies though, especially in the cases where these responses are not well understood.

Another relevant factor that was only briefly mentioned in [5] is taxis or advection, i.e. directed insect individual movement instead of (or additionally to) diffusive random motion. Directed movement may arise from an insect's behavioural responses to smell or light, or (for airborne species) can be a result of transport by the wind. Whilst the latter was shown to be

important on the large (landscape) scale, resulting in the synchronization of population oscillations [5,14], the effect of taxis is likely to be important on the small (single trap) and intermediate (single field) spatial scales. As was pointed out by Luiz Rodrigues in his commentary [2], on the single field scale, many insect species are known to respond to gradients of smell or odour [15] or, more generally, to the gradient in density of a resource [16,17]. Their directed movement along or against this gradient may then lead to pattern formation because of the so-called convective instability [17,18]. The population aggregation emerging due to pattern formation may significantly decrease the accuracy of population size estimation [5,19,20]. On the single trap scale, good understanding of taxis becomes especially important in the case of baited traps (e.g. pheromone-baited or light-baited), that are widely used in practice, and this should indeed become a direction of future research. We mention here that modelling of trap counts by baited traps is more complicated compared to the pitfall traps considered in our review as the effect of the attractant may change insect movement behaviour in a complex way [21,22]. Also, the currently available biological data about the possible effect of an attractant on the insect's movement behaviour are often ambiguous and insufficient.

The need for efficient and reliable pest insect monitoring as a crucial component of IPM was further emphasized by Bernd Blasius [3] and Edward Codling [4] who also pointed out the importance of identification of the physical and biological processes involved. Consequently, this raises the subtle and sometimes controversial issue of an adequate mathematical framework to describe these processes. However, what are adequate mathematical models and/or techniques is not always clear and the whole notion of adequacy may depend on the benchmark information available [23]. Once the process has been identified, usually it is not difficult to build a mathematical model for it, though the model might not necessarily be unique, as different mathematical tools may sometimes be used. For instance, the same movement pattern can be described by a partial differential equation (i.e. the diffusion equation) and by an integro-difference equation (i.e. the master equation); e.g. see [5], Section 2.2. The task of choosing 'the right model' becomes a much bigger challenge when we know the pattern but only part of the information about the actual process behind it is available; the choice of an adequate model then may become context-specific.

Identification of the movement pattern and the choice of the model, in particular as it applies to individual insect movement, was the main point of the commentary by Edward Codling [4]. A situation similar to that outlined in the previous paragraph can arise in the analysis of trap counts collected by a single trap. It appears that the same time course of counts is generated by a population of insects performing Lévy flight (LF), and by insects performing Brownian motion (BM) with a time-dependent diffusivity; see [5], Section 2.5. LF is generally regarded as a movement pattern basically different from BM, yet when considered in the particular context of insect trapping, they appear to be equivalent. Surprising as it may seem, it evokes other recent results showing that, in many cases and for many different reasons, the pattern produced by an animal population performing BM may become indistinguishable from the pattern produced by animals performing LF [24-26]. Altogether, this seems to lead to a paradigm shift: it is not the movement pattern itself that matters but the ecological context where the movement is considered. One relevant context is given by an animal's behaviour, which is likely to become a core component of future movement studies. We also mention here that, although diffusive insect movement with time-dependent diffusivity has been observed in some field studies [17], it can also be used formally as a convenient mathematical framework to approximate Lévy flights [5].

Admittedly, our review leaves many questions unanswered. Multiscale pest monitoring (as well as, in fact, multiscale agroecology as a whole) is a new research field at its infancy. A lot of theoretical and experimental research is needed before it will become a fully developed branch of ecological sciences with well understood powers and a clearly defined range of applications. There is a long road ahead; however, there are hardly any alternatives. The empirical approaches currently employed in pest monitoring have been useful but, without a strong theoretical background, they have limited powers and, what is worse, their limits are unknown. The ongoing global climate change exacerbates this by increasing the uncertainty of prediction. The 21st century needs efficient and environmentally-friendly agriculture that is built on theory-based knowledge and modern information technologies. We hope that the recent research reviewed in [5] may help to eventually reach this ambitious goal.

Acknowledgements

This work was supported by The Leverhulme Trust through grant F/00-568/X.

References

- [1] Tyson R. Pest control: A modelling approach. Comment on “Multiscale approach to pest insect monitoring: Random walks, pattern formation, synchronization, and networks” by S. Petrovskii, N. Petrovskaya, D. Bearup. *Phys Life Rev* 2014, this issue.
- [2] Rodrigues L.A.D. Diffusion models: still promoting progresses. Comment on “Multiscale approach to pest insect monitoring: Random walks, pattern formation, synchronization, and networks” by S. Petrovskii, N. Petrovskaya, D. Bearup. *Phys Life Rev* 2014, this issue.
- [3] Blasius B. Make your trappings count: the mathematics of pest insect monitoring. Comment on “Multiscale approach to pest insect monitoring: Random walks, pattern formation, synchronization, and networks” by S. Petrovskii, N. Petrovskaya, D. Bearup. *Phys Life Rev* 2014, this issue.
- [4] Codling E. Pest insect movement and dispersal as an example of applied movement ecology. Comment on “Multiscale approach to pest insect monitoring: Random walks, pattern formation, synchronization, and networks” by S. Petrovskii, N. Petrovskaya, D. Bearup. *Phys Life Rev* 2014, this issue.
- [5] Petrovskii S, Petrovskaya N, Bearup D. Multiscale approach to pest insect monitoring: random walks, pattern formation, synchronization, and networks. *Phys Life Rev* 2014. <http://dx.doi.org/10.1016/j.plrev.2014.02.001> [this issue].
- [6] Levin SA. The problem of pattern and scale in Ecology. *Ecology* 1992; 73(6):1943-67.
- [7] May RM. The effects of spatial scale on ecological questions and answers. In: Edwards PJ, May RM, Webb NR, editors. *Large-Scale Ecology and Conservation Biology* (British Ecological Society Symposium Volumes Series 35). Oxford: Wiley-Blackwell; 1994. p. 1-18.
- [8] Hastings A, Petrovskii SV, Morozov A. Spatial ecology across scales. *Biol Lett* 2011; 7:163-5.

- [9] Lewis M, Maini P, Petrovskii SV. Dispersal, Individual Movement, and Spatial Ecology: A Mathematical Perspective. Springer Lecture Notes in Mathematics Vol. 2071. Berlin: Springer; 2013.
- [10] Haghani M, Fathipour Y, Talebi AA, Baniaméri V. Thermal requirement and development of *Liriomyza sativae* (Diptera: Agromyzidae) on cucumber. *J Econ Entomol* 2007; 100:350-6.
- [11] Hou Y, Weng Z. Temperature-dependent development and life table parameters of *Octodonta nipae* (Coleoptera: Chrysomelidae). *Envir Entomol* 2010; 39:1676-84.
- [12] Doerr MD, Brunner JF, Jones VP. Phenology of *Lacanobia subjuncta* (Lepidoptera: Noctuidae) in Washington and Oregon apple orchards. *J Econom Entomol* 2005; 98:2100-6.
- [13] Jones VP, Hilton R, Brunner JF, Bentley WJ, Alston DG, Barrett B, et al. Predicting the emergence of the codling moth, *Cydia pomonella* (Lepidoptera: Tortricidae), on a degree-day scale in North America. *Pest Management Sci* 2013; 69(12):1393-8.
- [14] Bearup D, Petrovskii SV, Blackshaw R, Hastings A. Synchronized dynamics of *Tipula paludosa* metapopulation in a South-Western Scotland agroecosystem: linking pattern to process. *Am Nat* 2013; 182:393-409.
- [15] Rodrigues LAD, Varriale MC, Godoy WAC, Mistro DC. Spatiotemporal dynamics of an insect population in response to chemical substances. *Ecol Complex* 2013; 16:51-8.
- [16] Keller EF, Segel LA. Initiation of slime mold aggregation viewed as instability. *J Theor Biol* 1970; 26:399-415.
- [17] Okubo A, Levin SA. Diffusion and Ecological Problems: Modern Perspectives. Berlin: Springer; 2001.
- [18] Sherratt JA, Dagbovie AS, Hilker FM. A mathematical biologist's guide to absolute and convective instability. *Bull Math Biol* 2014; 76:1-26.
- [19] Petrovskaya NB, Petrovskii SV. The coarse-grid problem in ecological monitoring. *Proc R Soc A* 2010; 466:2933-53.
- [20] Petrovskaya NB, Petrovskii SV, Murchie AK. Challenges of ecological monitoring: estimating population abundance from sparse trap counts. *J R Soc Interface* 2012; 9:420-35.
- [21] Yamanaka T, Tatsuki S, Shimada M. An individual-based model for sex-pheromone-oriented flight patterns of male moths in a local area. *Ecol Mod* 2003; 161:35-51.
- [22] Bearup D, Petrovskii SV, Benefer C, Blackshaw R. Light trap count dynamics modelled by advection-diffusion equations. In preparation.
- [23] Petrovskii SV, Petrovskaya NB. Computational ecology as an emerging science. *Interface Focus* 2012; 2:241-254.

- [24] Plank MJ, Codling, EA. Sampling rate and misidentification of Lévy and non-Lévy movement paths. *Ecology* 2009; 90:3546-53.
- [25] Edwards AM. Overturning conclusions of Lévy flight movement patterns by fishing boats and foraging animals. *Ecology* 2011; 92:1247-57.
- [26] Petrovskii SV, Mashanova A, Jansen VAA. Variation in individual walking behavior creates the impression of a Lévy flight. *Proc Natl Acad Sci USA* 2011; 108: 8704-7.